

# Activities After Hip Replacement

After having a hip replacement, you may expect your lifestyle to be a lot like how it was before surgery—but without the pain. In many ways, you are right, but returning to your everyday activities will take time. Being an active participant in the healing process can help you get there sooner and ensure a more successful outcome.

Even though you will be able to resume most activities, you may have to change the way you do them. For example, you may have to learn new ways of bending down that keep your new hip safe. The suggestions you find here will help you enjoy your new hip while you safely resume your daily routines.

## Hospital Discharge

Your hospital stay will typically last from 1 to 4 days, depending on the speed of your recovery. If your hip replacement is done on an outpatient basis, you will go home on the same day as surgery.

Before you are discharged from the hospital, you will need to accomplish several goals, such as:

- Getting in and out of bed by yourself.
- Having acceptable pain control.
- Being able to eat, drink, and use the bathroom.
- Walking with an assistive device (a cane, walker, or crutches) on a level surface and being able to climb up and down two or three stairs.
- Being able to perform the prescribed home exercises.
- Understanding any hip precautions you may have been given to prevent injury and ensure proper healing.

If you are not yet able to accomplish these goals, it may be unsafe for you to go directly home after discharge. If this is the case, you may be temporarily transferred to a rehabilitation or skilled nursing center.

When you are discharged, your healthcare team will provide you with information to support your recovery at home. Although the complication rate after total hip replacement is low, when complications occur they can prolong or limit full recovery. Hospital staff will discuss possible complications, and review with you the warning signs of an infection or a blood clot.

## ***Warning Signs of Infection***

- Persistent fever (higher than 100 degrees)
- Shaking chills
- Increasing redness, tenderness or swelling of your wound
- Drainage from your wound
- Increasing pain with both activity and rest

## ***Warning Signs of a Blood Clot***

- Pain in your leg or calf unrelated to your incision
- Tenderness or redness above or below your knee
- Severe swelling of your thigh, calf, ankle or foot

In very rare cases, a blood clot may travel to your lungs and become life-threatening. Signs that a blood clot has traveled to your lungs include:

- Shortness of breath
- Sudden onset of chest pain
- Localized chest pain with coughing

Notify your doctor immediately if you develop any of the above signs.

## **Recovery at Home**

You will need some help at home for anywhere from several days to several weeks after discharge. Before your surgery, arrange for a friend, family member or caregiver to provide help at home.

### ***Preparing Your Home***

The following tips can help make your return home more comfortable, and can be addressed before your surgery:

- Rearrange furniture so you can maneuver with a cane, walker, or crutches. You may temporarily change rooms (make the living room your bedroom, for example) to minimize the use of stairs.
- Place items you use frequently (phone, remote control, glasses, pitcher and glass, reading material and medications, for example) within easy reach so you do not have to reach up or bend down.



Prepare a "recovery center" by placing items that you use frequently within easy reach.

- Remove any throw rugs or area rugs that could cause you to slip. Securely fasten electrical cords around the perimeter of the room.
- Get a good chair—one that is firm and has a higher-than-average seat. This type of chair is safer and more comfortable than a low, soft-cushioned chair.
- Install a shower chair, gripping bar, and raised toilet seat in the bathroom.
- Use assistive devices such as a long-handled shoehorn, a long-handled sponge, and a grabbing tool or reacher to avoid bending over too far.

## ***Wound Care***

During your recovery at home, follow these guidelines to take care of your wound and help prevent infection:

- Keep the wound area clean and dry. A dressing will be applied in the hospital and should be changed as often as directed by your doctor. Ask for instructions on how to change the dressing before you leave the hospital.
- Follow your doctor's instructions on how long to wait before you shower or bathe.
- Notify your doctor immediately if the wound appears red or begins to drain. This could be a sign of infection.

## ***Swelling***

Expect moderate to severe swelling in the first few weeks after surgery. You may also have mild to moderate swelling for 3 to 6 months after surgery. To reduce swelling, elevate your leg slightly and apply ice. Wearing compression stockings may also help reduce swelling. Notify your doctor if you experience new or severe swelling, since this may be the warning sign of a blood clot.

## ***Medication***

Take all medications as directed by your doctor. Home medications may include opioid and non-opioid pain pills, oral or injectable blood thinners, stool softeners, and anti-nausea medications.

Be sure to talk with your doctor about all your medications—even over-the-counter drugs, supplements, and vitamins. Your doctor will tell you which over-the-counter medicines are safe to take while using prescription pain medication.

It is especially important to prevent any bacterial infections from developing in your artificial joint. Some patients with special circumstances may be required to take antibiotics prior to dental work to help prevent infection. Ask your doctor if you should take antibiotics before dental work. You may also wish to carry a medical alert card so that, if an emergency arises, medical personnel will know that you have an artificial joint.

## ***Diet***

By the time you go home from the hospital, you should be eating a normal diet. Your doctor may recommend that you take iron and vitamin supplements. You also may be advised to avoid supplements that include vitamin K and foods rich in vitamin K if you are taking the blood thinner medication warfarin (Coumadin). Foods rich in vitamin K include broccoli, cauliflower, brussel sprouts, liver, green beans, garbanzo beans, lentils, soybeans, soybean oil, spinach, kale, lettuce, turnip greens, cabbage, and onions.

Continue to drink plenty of fluids and avoid alcohol. You should continue to watch your weight to avoid putting more stress on the joint.

## **Resuming Normal Activities**

Once you get home, you should stay active. The key is to not do too much, too soon. While you can expect some good days and some bad days, you should notice a gradual improvement over time. Generally, the following guidelines will apply:

### ***Weight Bearing***

Follow your doctor's specific instructions about the use of a cane, walker, or crutches and when you can put weight on the leg. Full weight bearing may be allowed immediately or may be delayed by several weeks depending on the type of hip replacement you have undergone and your doctor's instructions.

### ***Driving***

In most cases, it is safe to resume driving when you are no longer taking opioid pain medication, and when your strength and reflexes have returned to a more normal state. Your doctor will help you determine when it is safe to resume driving.

## ***Sexual Activity***

Please consult your doctor about how soon you can safely resume sexual activity. Depending on your condition, you may be able to resume sexual activity within several weeks after surgery.

## ***Sleeping Positions***

Depending on your surgery, your doctor may ask you to avoid certain sleeping positions or to sleep with a pillow between your legs for a length of time. Ask your doctor which sleeping positions are safest and most appropriate for you.

## ***Return to Work***

Depending on the type of activities you do on the job and the speed of your recovery, it may take from several days to several weeks before you are able to return to work. Your doctor will advise you when it is safe to resume your normal work activities.

## ***Sports and Exercise***

Continue to do the exercises prescribed by your physical therapist for at least 2 months after surgery. In some cases, your doctor may recommend riding a stationary bicycle to help maintain muscle tone and keep your hip flexible.

As soon as your doctor gives you the go-ahead, you can return to many of the sports activities you enjoyed before your hip replacement:

- Walk as much as you would like, but remember that walking is no substitute for the exercises prescribed by your doctor and physical therapist.
- Swimming is an excellent low-impact activity after a total hip replacement; you can begin swimming as soon as the wound is sufficiently healed. Your doctor will let you know when you can begin.
- In general, lower impact fitness activities such as golfing, bicycling, and light tennis, put less stress on your hip joint and are preferable over high-impact activities such as jogging, racquetball and skiing.

## ***Air Travel***

Pressure changes and immobility may cause your operated leg to swell, especially if it is just healing. Ask your doctor before you travel on an airplane. When going through security, be aware that the sensitivity of metal detectors varies and your artificial joint may cause an alarm. Tell the screener about your artificial joint before going through the metal detector.

## **Dos and Don'ts to Protect Your New Hip**


Dos and don'ts (precautions) vary depending on your doctor's surgical technique and preferences. Your doctor and physical therapist will provide you with a list of dos and don'ts to remember with your new hip. These precautions will help to prevent the new joint from dislocating and ensure proper healing. Some of the most common precautions are listed below. Ask your doctor if these precautions apply to you.

### ***The Don'ts***

- Don't cross your legs at the knees for at least 6 to 8 weeks.
- Don't bring your knee up higher than your hip.
- Don't lean forward while sitting or as you sit down.
- Don't try to pick up something on the floor while you are sitting.
- Don't turn your feet excessively inward or outward when you bend down.
- Don't reach down to pull up blankets when lying in bed.
- Don't bend at the waist beyond 90 degrees.

### ***The Dos***

- Do keep the leg facing forward.
- Do keep the affected leg in front as you sit or stand.
- Do use a high kitchen or barstool in the kitchen.
- Do kneel on the knee on the operated leg (the bad side).
- Do use ice to reduce pain and swelling, but remember that ice will diminish sensation. Don't apply ice directly to the skin; use an ice pack or wrap it in a damp towel.
- Do apply heat before exercising to assist with range of motion. Use a heating pad or hot, damp towel for 15 to 20 minutes.
- Do cut back on your exercises if your muscles begin to ache, but don't stop doing them!



Information on this topic is also available as an **OrthoInfo Basics PDF Handout**.

For more information:

### **Basics Handouts**

[\(/en/about-orthoinfo/OrthoInfo-Basics/basics-handouts/\)](/en/about-orthoinfo/OrthoInfo-Basics/basics-handouts/)

### **Last Reviewed**

December 2018

### **Contributed and/or Updated by**

[Jared R.H. Foran, MD](#)

### **Peer-Reviewed by**

[Stuart J. Fischer, MD](#)

AAOS does not endorse any treatments, procedures, products, or physicians referenced herein. This information is provided as an educational service and is not intended to serve as medical advice. Anyone seeking specific orthopaedic advice or assistance should consult his or her orthopaedic surgeon, or locate one in your area through the AAOS [Find an Orthopaedist](#) program on this website.